

This quiz has 2 questions.

1. (10 points) Find the eigenvalues and eigenvectors of the following matrix.

$$\mathbf{A} = \begin{bmatrix} 2 & -1 & 1 \\ 1 & 0 & 1 \\ 2 & -2 & 3 \end{bmatrix}.$$

$$A - \lambda I = \begin{bmatrix} 2-\lambda & -1 & 1 \\ 1 & -\lambda & 1 \\ 2 & -2 & 3-\lambda \end{bmatrix}, \quad \det(A - \lambda I) = (2-\lambda)(-\lambda)(3-\lambda) - 2 -$$

$$-2 + 2\lambda - 2(2-\lambda) + (3-\lambda) = (2-\lambda)(-\lambda)(3-\lambda) + (3-\lambda) =$$

$$= (3-\lambda)(\lambda^2 - 2\lambda + 1) = (3-\lambda)(1-\lambda)^2$$

$$\lambda_1 = 3, \quad \lambda_2 = 1 \leftarrow \text{double eigenvalue.}$$

$$\text{For } \lambda_1 = 3 \quad A - 3I = \begin{bmatrix} -1 & -1 & 1 \\ 1 & -3 & 1 \\ 2 & -2 & 0 \end{bmatrix} \xrightarrow{\substack{R2+R1 \\ R3+2R1}} \begin{bmatrix} -1 & -1 & 1 \\ 0 & -4 & 2 \\ 0 & -4 & 2 \end{bmatrix} \xrightarrow{R3-R2} \begin{bmatrix} -1 & -1 & 1 \\ 0 & -4 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

$$v_1 = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \quad -4y + 2z = 0, \quad y = \frac{z}{2}, \quad -x - y + z = 0, \quad -x - \frac{z}{2} + z = 0$$

$$x = \frac{z}{2} \quad \begin{pmatrix} \frac{z}{2} \\ \frac{z}{2} \\ z \end{pmatrix} = z \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ 1 \end{pmatrix} \quad v_1 = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \\ 1 \end{pmatrix}.$$

$$\text{For } \lambda = 1 \quad A - I = \begin{bmatrix} 1 & -1 & 1 \\ 1 & -1 & 1 \\ 2 & -2 & 2 \end{bmatrix} \xrightarrow{\substack{R2-R1 \\ R3-2R1}} \begin{bmatrix} 1 & -1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$v_2 = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$

$$x - y + z = 0, \quad x = y - z, \quad \begin{pmatrix} y - z \\ y \\ z \end{pmatrix} = \begin{pmatrix} y \\ y \\ 0 \end{pmatrix} + \begin{pmatrix} -z \\ 0 \\ z \end{pmatrix} = y \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} + z \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}.$$

$$\text{Thus } v_2 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \text{ and } v_3 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}.$$

2. (10 points) For matrix \mathbf{A} from problem 1, which of the eigenvalues are complete? Is matrix \mathbf{A} complete? Is \mathbf{A} diagonalizable? If so, find matrices \mathbf{A} and Λ such that $\mathbf{A} = \mathbf{S}\Lambda\mathbf{S}^{-1}$. Explain your answers.